



Cambridge Assessment International Education
Cambridge Ordinary Level

CHEMISTRY

5070/42

Paper 4 Alternative to Practical

May/June 2019

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **8** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).


GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Question	Answer	Marks																				
1(a)	A = conical flask / Erlenmeyer flask (1) B = pipette (1) C = burette (1)	3																				
1(b)(i)	<table border="1" data-bbox="322 416 1077 938"> <thead> <tr> <th data-bbox="322 416 551 517">titration number</th> <th data-bbox="551 416 680 517">1</th> <th data-bbox="680 416 810 517">2</th> <th data-bbox="810 416 940 517">3</th> <th data-bbox="940 416 1077 517">4</th> </tr> </thead> <tbody> <tr> <td data-bbox="322 517 551 617">final burette reading / cm³</td> <td data-bbox="551 517 680 617">24.1</td> <td data-bbox="680 517 810 617">47.5</td> <td data-bbox="810 517 940 617">23.6</td> <td data-bbox="940 517 1077 617">24.4</td> </tr> <tr> <td data-bbox="322 617 551 735">initial burette reading / cm³</td> <td data-bbox="551 617 680 735">0.0</td> <td data-bbox="680 617 810 735">23.7</td> <td data-bbox="810 617 940 735">0.3</td> <td data-bbox="940 617 1077 735">0.8</td> </tr> <tr> <td data-bbox="322 735 551 938">volume of 0.600 mol / dm³ hydrochloric acid used / cm³</td> <td data-bbox="551 735 680 938">24.1</td> <td data-bbox="680 735 810 938">23.8</td> <td data-bbox="810 735 940 938">23.3</td> <td data-bbox="940 735 1077 938">23.6</td> </tr> </tbody> </table>	titration number	1	2	3	4	final burette reading / cm ³	24.1	47.5	23.6	24.4	initial burette reading / cm ³	0.0	23.7	0.3	0.8	volume of 0.600 mol / dm ³ hydrochloric acid used / cm ³	24.1	23.8	23.3	23.6	3
titration number	1	2	3	4																		
final burette reading / cm ³	24.1	47.5	23.6	24.4																		
initial burette reading / cm ³	0.0	23.7	0.3	0.8																		
volume of 0.600 mol / dm ³ hydrochloric acid used / cm ³	24.1	23.8	23.3	23.6																		
1(b)(ii)	23.7 (cm ³)	1																				
1(b)(iii)	(water) only changes concentration / does not affect amount or moles (of Mg(OH) ₂)	1																				
1(c)(i)	0.01398 / 1.398 × 10 ⁻²	1																				
1(c)(ii)	0.00699 / 6.99 × 10 ⁻³	1																				
1(c)(iii)	M1 M _r = 58 (1) M2 0.405 (g) (1)	2																				
1(c)(iv)	11.7 (g)	1																				
1(c)(v)	3.47(%)	1																				

Question	Answer	Marks
2	<p>MARKS CAN SCORE IN DIAGRAM AND IN TEXT</p> <p>M1 chromatography methods M2 spots of sample and E150a and E155 (1) M3 solvent in jar / beaker with level below base line (or on diagram) (1) M4 baseline drawn in pencil (1) M5 let solvent run (to near top of paper) OR solvent front mentioned (1) M6 compare position of spot(s) from sample with spots from E150a / E155 OR (measure) Rf / retention factor (1)</p>	6

Question	Answer	Marks
3(a)(i)	<p>M1 cathode: pink / brown and solid M2 anode: bubbles / fizzing / effervescence</p>	2
3(a)(ii)	<p>M1 glowing splint (into tube of gas) M2 relights / rekindles</p>	2
3(a)(iii)	<p>M1 colour fades / becomes paler blue / goes colourless M2 as Cu^{2+} / copper ions concentration falls</p>	2
3(b)(i)	colour stays the same	1
3(b)(ii)	<p>M1 mass of cathode increases (1) M2 mass of anode decreases (1) M3 copper removed from anode and copper forms on cathode (1)</p>	3
3(b)(iii)	copper purification / copper refining / electroplating	1

Question	Answer	Marks
4(a)	solution turns brown	1
4(b)(i)	(calcium carbonate) neutralises the acid	1
4(b)(ii)	increase rate (of reaction)	1
4(b)(iii)	effervescence / bubbles / fizzing OR (calcium carbonate) dissolves / disappears	1
4(c)(i)	<p>POINTS</p>  <p>6 / 7 points correct = 2 marks 4 / 5 points correct = 1 mark</p>	2
4(c)(ii)	ANOMALOUS POINT mark is for circle round point at 100, 11	1
4(c)(iii)	CURVE smooth curve through all the points except the anomalous one	1
4(c)(iv)	13.5 (cm ³)	1
4(c)(v)	80 (s)	1

Question	Answer	Marks
4(c)(vi)	M1 gradient or slope decreases (1) M2 rate decreases (with time) (1) M3 as concentration decreases (1) M4 lower collision frequency (1)	4

Question	Answer	Marks
5(a)	M1 add copper oxide or copper carbonate or copper hydroxide (1) M2 dilute sulfuric acid (1) M3 (solid added until in) excess / no more dissolves / solid is seen / no more bubbles (with carbonate) M4 filter (the solution to remove excess solid) M5 heat / warm / evaporate / boil / leave (the solution) AND until signs of crystal formation / saturated (solution) / crystallisation (point) / crystals form on glass rod or other surface / crystals can be seen	5
5(b)	M1 (mix)silver nitrate solution / aqueous silver nitrate M2 with any named solution of a soluble chloride M3 filter M4 wash (residue) with water and dry on filter paper	4

Question	Answer	Marks
6(a)	M1 carbon dioxide (1) M2 carbonate present (1)	2
6(b)	white precipitate	1
6(c)	M1 white precipitate (1) M2 insoluble in excess (1)	2
6(d)	no (visible) change / no reaction / no precipitate seen / slight (white) precipitate	1