



Cambridge International Examinations
Cambridge Ordinary Level

PHYSICS

5054/21

Paper 2 Theory

October/November 2016

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2016 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2016	5054	21

Section A

- 1 (a) velocity/it has a direction/is a vector B1
- (b) (i) $(F =)ma$ or 800×1.5 C1
1200 N A1
- (ii) friction/air resistance acts on car B1
opposes force due to engine B1
- (iii) $(\Delta v =)at$ or 1.5×4.0 or 6.0 C1
31 m/s A1 [7]
- 2 (a) 260 N B1
- (b) (i) for a body in equilibrium B1
(total) clockwise moment = (total) anticlockwise moment B1
- (ii) $F_1d_1 = F_2d_2$ or 260×0.35 or 91 or $F \times 0.65$ C1
 $260 \times 0.35 = F \times 0.65$ or $260 \times 0.35/0.65$ or $91 = F \times 0.65$ or $91/0.65$ C1
140 N A1 [6]
- 3 (a) chemical (potential energy) B1
- (b) (i) non-renewable **and** oil/it is not replaced/will run out B1
- (ii) acid rain **or** produces CO₂ **or** warms lakes/rivers/sea **or** global warming **or** greenhouse effect B1
- (c) (i) useful energy output/ (total) energy input **or** power for energy twice B1
- (ii) 1 $1.9 \times 10^9/0.38$ or $1.9 \times 10^9 \times 100/38$ C1
 5.0×10^9 W A1
- 2 $(E =)Pt$ or $0.62 \times 5.0 \times 10^9 \times 2.0 (\times 3600)$ or $(5.0 - 1.9) \times 10^9$ etc. C1
 2.2×10^{13} J A1 [8]
- 4 (a) smallest angle for total internal reflection **or** angle for refraction along surface B1
angle of incidence in (optically) denser medium B1
- (b) vertical ray continues undeviated B1
second ray (60° to horizontal) refracts away from normal into the air B1
third ray reflects internally **and** $i = r$ by eye **not** if any refracted ray B1 [5]


Page 3	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2016	5054	21

- 5 (a) number of oscillations / vibrations / wavelengths / compressions / rarefactions / cycles per second / unit time B1
- (b) (i) ($\lambda =$) c/f or 330 / 2200 C1
0.15 m A1
- (ii) 1 no change
and
2 increases B1
- (c) (i) 1 loudspeaker vibrates / oscillates / moves to and fro (and collides with molecules) B1
2 compressions **and** rarefactions / molecules vibrate / longitudinal wave vibration / oscillation / energy passed on B1
B1
- (ii) fewer / no molecules / particles **and** less / no energy / vibration transferred B1 [8]
- 6 (a) (i) X N-pole B1
Y S-pole **and** Z N-pole B1
- (ii) they touch / move towards each other **and** opposite poles attract B1
- (b) any sensible use: starting-motor circuit; with a logic gate; nuclear power station B1
corresponding explanation: current too large for dash-board switch; current too small to power device; too dangerous to reach switch B1 [5]
- 7 (a) (i) supplies the (mains) e.m.f. / voltage B1
- (ii) to complete the circuit / is at 0 V B1
- (b) (i) the circuit / supply is cut / broken or current stops B1
fuse melts / blows / burns B1
- (ii) live wire B1
when it cuts the circuit / melts no part of the appliance is live / no shock B1 [6]

[45]

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2016	5054	21

Section B

- 8 (a) (i) 11 protons **and** 11 electrons B1
 13 neutrons B1
 electrons in orbit/surrounding nucleus **or** neutrons **and** protons in nucleus B1
- (ii) **one** more neutron (in sodium-24) **or** one fewer neutron in sodium-23 B1 [4]
- (b) (i) electron B1
- (ii) ${}_{-1}^0(\beta)$ **cao** B1
 ${}^{24}(\text{Mg})$ B1
 ${}_{12}(\text{Mg})$ B1 [4]
- (c) electromagnetic (radiation/rays/waves) M1
 (very) high frequency/energy **or** (very) short wavelength A1 [2]
- (d) (i) path curving upwards B1
 (ii) path continues in straight line B1
 (iii) beta-particle charged **or** gamma-ray uncharged B1 [3]
- (e) long enough **or** short enough B1
 to take measurements **or** so the body is not irradiated for long B1 [2]
[15]
- 9 (a) (i) magnetic field mentioned B1
 alternating/changing magnetic field B1
 current/voltage/e.m.f. induced (in secondary coil) B1
- (ii)  B1
 diode B1 [5]
- (b) (i) work done/energy transferred per unit charge M1
 electrical energy to other forms **or** for whole circuit **or** property of supply A1

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2016	5054	21

(ii)	1	1.3V	B1	
	2	$(I =)V/R$ or 1.3/5.2 0.25 A	C1 A1	
	3	$(Q =)It$ or $0.25 \times 1.5 \times 3600$ or 0.25×1.5 $0.25 \times 1.5 \times 3600$ or 0.37/0.375/0.38 1300/1350/1400 C	C1 C1 A1	[8]
(c)		plastic/casing is an (electrical) insulator no shock possible	M1 A1	[2] [15]
10 (a)	(i)	molecules / they close together or small gaps between molecules	B1	
	(ii)	molecules / they exert large (repulsive) forces	B1	[2]
(b)	(i)	$(V =)m/\rho$ or 680/0.85 800 cm^3 or $8.0 \times 10^{-4} \text{ m}^3$	C1 A1	
	(ii)	1 molecules <u>vibrate</u> collide with neighbours transfer vibration / energy	B1 B1 B1	
		2 heated / hot oil expands / less dense rises convection current/circulation established	B1 B1 B1	
		3 any suitable named insulator and it is a poor conductor	B1	[9]
(c)	(i)	<u>temperature</u> at which (liquid) vaporises / becomes gas / steam	B1	
	(ii)	$(Q =)mc \Delta T$ or $680 \times 2.0 \times (260 - 20)$ or $680 \times 2.0 \times 240$ $3.3 \times 10^5 \text{ J}$	C1 A1	
	(iii)	heat supplied to <u>pan</u> or heat lost to <u>air / surroundings</u>	B1	[4] [15]