

# Mark Scheme (Results)

## Summer 2015

Pearson Edexcel International GCSE in  
Chemistry (4CH0) Paper 1CR

Pearson Edexcel International in Science  
Double Award (4SC0) Paper 1CR

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2015

Publications Code UG041121

All the material in this publication is copyright

© Pearson Education Ltd 2015

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a)	Diagram shows four circles well-spaced apart	accept minimum of 3 complete circles ignore size and shape of circles ignore arrows and other symbols implying movement  ignore a pattern reject any touching circles reject circles joined by bonds no penalty for half-circles at edges of square	1
(b)	move freely/randomly	Accept fast OWTTE ignore references to vibrate	1
(c)	<b>M1</b> – (average kinetic) energy of the particles increases  <b>M2</b> – <u>more</u> particles have enough energy to escape / particles can escape <u>more</u> easily OR <u>more</u> particles overcome the forces (of attraction) holding them together (in the liquid) OR the forces (of attraction) between the particles are overcome <u>more</u> often	accept particles move faster/more rapidly/more quickly allow the energy of the liquid increases  accept particles escape <u>more</u> quickly  accept molecules/atoms for particles for both M1 and M2  allow bonds for force of attraction	2
<b>Total 4 marks</b>			

Question number	Answer	Notes	Marks								
2 (a)	<u>fractional</u> distillation	accept fractionation	1								
(b)	<table border="1" data-bbox="394 344 1055 624"> <thead> <tr> <th data-bbox="394 344 629 416">Fraction</th> <th data-bbox="629 344 1055 416">Description</th> </tr> </thead> <tbody> <tr> <td data-bbox="394 416 629 488">A</td> <td data-bbox="629 416 1055 488">it contains only gases</td> </tr> <tr> <td data-bbox="394 488 629 560">F</td> <td data-bbox="629 488 1055 560">it is the most viscous</td> </tr> <tr> <td data-bbox="394 560 629 624">F</td> <td data-bbox="629 560 1055 624">it contains bitumen</td> </tr> </tbody> </table>	Fraction	Description	A	it contains only gases	F	it is the most viscous	F	it contains bitumen		<p>1</p> <p>1</p> <p>1</p>
Fraction	Description										
A	it contains only gases										
F	it is the most viscous										
F	it contains bitumen										
(c)	as the number of carbon atoms/it/they increases the boiling point increases	<p>accept reverse argument</p> <p>allow positive correlation</p> <p>ignore (directly) proportional</p> <p>ignore references to hydrogen atoms</p>	1								
<b>Total 5 marks</b>											

Question number	Answer	Notes	Marks
3 (a)	<p><b>M1</b> – C</p> <p><b>M2</b> – (it) has a spot in line with/at the same height as (the spot produced by) bute/an illegal drug</p>	<p>Accept references to travelling same distance / having same <math>R_f</math> value</p> <p><b>M2</b> dep on <b>M1</b></p>	<p>1</p> <p>1</p>
(b)	a substance/liquid that dissolves a solute/solid/another substance	Accept it forms a solution with a solute/solid/substance	1
(c)	<p><b>M1</b> <math>\frac{\text{correctly measured distance for lasix spot}}{\text{correctly measured distance of solvent front}}</math></p> <p><b>M2</b> – any value in range 0.73 – 0.77</p>	<p>Lasix spot 62-64 mm / 6.2-6.4 cm Solvent front 84 mm / 8.4 cm</p> <p>Minimum of 2 dp correct answer with no working scores 2</p> <p><b>M2</b> csq on <b>M1</b></p>	<p>1</p> <p>1</p>
(d)	the more soluble the substance the further it will travel	Allow distance increases with (increasing) solubility ignore any reference to proportionality	1
<b>Total 6 marks</b>			

Question number	Answer		Notes	Marks
4 (a)	<b>Description of reaction</b>	<b>Metal</b>	3 correct = 2 marks 1 correct = 1 mark  accept symbols	2
	it explodes on contact with water	caesium		
	it fizzes gently	lithium		
	it reacts violently and forms a lilac flame	potassium		
(b) (i)	<b>M1</b> – hydrogen		ignore symbol or formula even if incorrect	1
	<b>M2</b> – H <sub>2</sub>		reject H accept H <sub>2</sub> (g) as a <u>product</u> in an equation  ignore name even if incorrect accept LiOH as a <u>product</u> in an equation	1
(ii)	<b>M1</b> – lithium hydroxide		ignore formula even if incorrect	1
	<b>M2</b> – LiOH		ignore name even if incorrect	1
(iii)	<b>M1</b> – add (red) litmus		accept any named indicator	1
	<b>M2</b> - turns blue		accept correct colour for named indicator	1
	OR		ignore purple	
	<b>M1</b> – use a pH meter / measure pH			
<b>M2</b> - pH > 7		M2 DEP on M1 do not award <b>M1</b> or <b>M2</b> if blue litmus is used		
<b>Total 8 marks</b>				

Question number	Answer	Notes	Marks
5 (a) (i)	<b>M1</b> – E		1
	<b>M2</b> – volume of carbon dioxide/gas (given off) is half / is 30 cm <sup>3</sup> (not 60 cm <sup>3</sup> )	accept volume of carbon dioxide/gas is less accept amount for volume ignore references to rate in (i)	1
(ii)	<b>M1</b> – C		1
	<b>M2</b> – curve levels off later / curve is less steep	Accept the reaction is slower /carbon dioxide/gas given off <u>more</u> slowly / takes <u>longer</u> for reaction to complete	1
(iii)	<b>M1</b> – B		1
	<b>M2</b> – curve levels off earlier / curve is <u>steeper</u>	Accept the reaction is faster / carbon dioxide/gas given off <u>more</u> quickly / takes <u>less</u> time for reaction to complete  ignore references to collision theory throughout part (a)  <b>M2</b> dep on <b>M1</b> correct or missing for all answers to part (a)	1
(b)	(Gas) syringe / measuring cylinder (over water) / burette (over water)	Allow graduated tube	1
<b>Total 7 marks</b>			





(v)	7 (cm <sup>3</sup> )	Ignore no reaction / reaction not started accept any value between 6.8 and 7.2	
<b>Total 9 marks</b>			

Question number	Answer	Notes	Marks
7 (a) (i)	(saturated) – <u>all</u> (carbon to carbon) bonds are single / no (carbon to carbon) double bonds	accept no (carbon to carbon) multiple bonds ignore any references to hydrogen	1
(ii)	<b>M1</b> - (compounds/substances/molecules) containing hydrogen and carbon (atoms/elements)  <b>M2</b> - only	reject atoms/elements/ions/mixture in place of compounds reject compounds/substances/molecules in place of atoms/elements  accept other terms with same meaning, e.g. solely, exclusively, just	1  1
(iii)	<b>C</b> (C <sub>5</sub> H <sub>12</sub> )	<b>M2</b> DEP on mention of hydrogen and carbon / C and H and no other element	1
(b) (i)	$C_8H_{18} + 12.5O_2 \rightarrow 8CO_2 + 9H_2O$  <b>M1</b> – all formulae correct <b>M2</b> – balanced using correct formulae	accept multiples	2
(ii)	carbon monoxide	If both name and formula given, mark name only accept correct formula	1

Question number	Answer	Notes	Marks
7 (c)	(provides an alternative pathway of) lower activation energy	Accept (molecules adsorb onto catalyst and covalent) bonds weakened	1
(i)	(provides an alternative pathway of) lower activation energy	Accept (molecules adsorb onto catalyst and covalent) bonds weakened	1
(ii)	silica/silicon dioxide/alumina/aluminium oxide	accept correct formulae accept aluminosilicate(s) accept zeolite(s) ignore silica oxide and alumina oxide If both name and formula given, mark name only	1
(iii)	C <sub>2</sub> H <sub>4</sub>	Accept structural or displayed formula	1
(iv)	ethene	accept ethylene	1

Question number	Answer	Notes	Marks
8 (a) (i)	<p><b>M1</b> &amp; <b>M2</b> – any two from:</p> <ul style="list-style-type: none"> <li>• does not melt/high melting point</li> <li>• does not colour the flame</li> <li>• inert/unreactive / does not burn/react with oxygen/air</li> </ul>	Ignore general physical properties of metals, eg boiling point	2
(ii)	to remove any substance that may affect the colour	ignore references to removing impurities Allow result/flame for colour	1
(iii)	difficult to see the colour produced by the substance (under test)	Accept flame not hot enough (to vaporise the sample) Accept the temperature is not high enough (to vaporise the sample) Allow flame is (already) coloured	1
(b) (i)	<p>(X) <b>M1</b> – sodium <b>M2</b> – chloride</p> <p>(Y) <b>M3</b> – lithium <b>M4</b> – sulfate</p>	Accept symbol in any formula	1 1
(ii)	iron(II) / Fe <sup>2+</sup> / Fe <sup>+2</sup> / Fe <sup>++</sup>	accept Li symbol and SO <sub>4</sub> in any formula accept strontium for <b>M3</b> accept ferrous ignore iron ion if both name and formula given both must be correct	1

(c)	M1 – add (dilute) acid		1
	M2 – test gas/bubbles/carbon dioxide with limewater	If incorrect gas mentioned, only M1 can be awarded	1
	M3 – limewater turns milky	M3 DEP on mention of gas	1




9 (c)	$\text{Cl}_2 + 2 \text{NaOH} \rightarrow \text{NaCl} + \text{NaClO} + \text{H}_2\text{O}$ <b>M1</b> – all formulae correct <b>M2</b> – balanced using correct formulae		2
-------	--	--	---



Question number	Answer	Notes	Marks
10 (a) (i)	$\text{Zn(s)} + 2 \text{HCl(aq)} \rightarrow \text{ZnCl}_2\text{(aq)} + \text{H}_2\text{(g)}$ <p><b>M1</b> – all formulae correct and equation balanced</p> <p><b>M2</b> – state symbols correct</p>	<p><b>M2</b> can be awarded for near misses on formulae, e.g. ZnCl and H</p> <p>accept upper case letters for state symbols</p>	2
(b)	<p><b>M1</b> bubbles/fizzing/effervescence</p> <p><b>M2</b> zinc/solid gets smaller/disappears</p>	<p>accept gas given off ignore hydrogen given off</p> <p>accept zinc/solid dissolves / (final) solution is <u>colourless</u> reject zinc melts and other Group 1 observations, eg floats / moves across surface</p> <p>Ignore references to heat and temperature change</p>	2

Question number	Answer	Notes	Marks												
10 (c) (i)	<table border="1" data-bbox="427 252 1218 485"> <thead> <tr> <th></th> <th>Experiment 1</th> <th>Experiment 2</th> </tr> </thead> <tbody> <tr> <td>Final burette reading in cm<sup>3</sup></td> <td>10.40</td> <td>22.70</td> </tr> <tr> <td>Initial burette reading in cm<sup>3</sup></td> <td>0.00</td> <td>1.90</td> </tr> <tr> <td>Volume of acid added in cm<sup>3</sup></td> <td>10.40</td> <td>20.80</td> </tr> </tbody> </table> <p data-bbox="421 560 1205 738"> <b>M1</b> – all four burette readings correct  <b>M2</b> – subtractions correct  <b>M3</b> – all six values in table given to 2 decimal places         </p> <p data-bbox="338 847 1263 954">           (ii) <b>M1</b> – (because) the volume/amount of acid required has doubled  <b>M2</b> – the concentration is half / 0.37 (mol dm<sup>-3</sup>)         </p> <p data-bbox="409 994 461 1023"><b>OR</b></p> <p data-bbox="409 1066 1155 1094"><b>M1</b> for use of an expression such as <math>V_1c_1 = V_2c_2</math></p> <p data-bbox="409 1137 1305 1241"><b>M2</b> for indicating how <math>c_2</math> can be calculated (e.g. because <math>V_1, c_1,</math> and <math>V_2</math> are known) / for an answer of 0.37 (mol dm<sup>-3</sup>)</p>		Experiment 1	Experiment 2	Final burette reading in cm <sup>3</sup>	10.40	22.70	Initial burette reading in cm <sup>3</sup>	0.00	1.90	Volume of acid added in cm <sup>3</sup>	10.40	20.80	<p data-bbox="1339 560 1854 627">Ignore trailing zeroes for <b>M1</b> and <b>M2</b></p> <p data-bbox="1339 667 1854 734"><b>M2</b> CSQ on burette readings given in table</p> <p data-bbox="1350 847 1641 882">Mark independently</p> <p data-bbox="1339 1066 1794 1133">accept either a calculation or a description</p>	<p data-bbox="1944 312 1966 341">3</p> <p data-bbox="1944 852 1966 880">1</p> <p data-bbox="1944 924 1966 952">1</p>
	Experiment 1	Experiment 2													
Final burette reading in cm <sup>3</sup>	10.40	22.70													
Initial burette reading in cm <sup>3</sup>	0.00	1.90													
Volume of acid added in cm <sup>3</sup>	10.40	20.80													

Question number	Answer	Notes	Marks
11 (a) (i)		ignore bond angles Ignore brackets and n Do not penalise FI	1
	(ii) <b>M1</b> – a long chain (molecule)	accept large molecule / macromolecule	1
	<b>M2</b> – formed when (many) small molecules/monomers join (together)	Accept react/bond/add/link for join	1
	(iii) poly(tetrafluoroethene)/poly(tetrafluoroethylene)	accept names without brackets Ignore minor spelling errors Ignore PTFE accept Teflon	1
(b)	<b>M1</b> (name) – ethene	accept ethylene	1
	<b>M2</b> (formula) – C <sub>2</sub> H <sub>4</sub>	reject structural or displayed formula Penalise inappropriate use of upper and lower case letters or numbers No penalty for correct answers on wrong lines	1

(c)	<p><b>M1</b> – (they) do not biodegrade</p> <p><b>M2</b> – (because) they are inert / do not react / are unreactive</p>	<p>accept not broken down by bacteria / microbes / decomposers / microorganisms / enzymes</p> <p>ignore do not react with any named chemical ignore references to bond strengths / bond breaking</p> <p>Mark independently</p>	<p>1</p> <p>1</p>
-----	---	--	-------------------

Question number	Answer	Notes	Marks
12 (a)	copper	ignore symbol reject copper(II) / copper(II) ions / Cu <sup>2+</sup>	1
(b)	zinc cannot displace itself	Accept zinc cannot react with zinc ions/zinc nitrate Accept the two metals involved have the same reactivity	1
(c)	aluminium zinc M copper  <b>M1</b> – aluminium at top <u>and</u> copper at bottom  <b>M2</b> – zinc above M	award <b>M2</b> irrespective of where zinc is placed in the list	2

(d) (i)	oxidation <u>and</u> reduction occur <b>OR</b> electron loss <u>and</u> electron gain occur <b>OR</b> oxidation number increase <u>and</u> decrease	reject references to oxygen  Accept electron transfer  Ignore species involved	1
(ii)	<b>M1</b> – Ag <sup>+</sup> /silver <u>ion</u> (s)  <b>M2</b> – it gains electron/is reduced <b>OR</b> it takes electrons from Mg/magnesium (atoms) <b>OR</b> its oxidation number decreases <b>OR</b> it causes the oxidation number of Mg to increase	<b>M2</b> DEP on <b>M1</b> or near miss, e.g. Ag	1  1

Question number	Answer	Notes	Marks
13 (a)	measuring cylinder/measuring jug	accept burette/pipette	1
(b)	no more bubbles/fizzing/effervescence/gas given off <b>OR</b> solid/zinc carbonate can be seen in the beaker <b>OR</b> the solid/zinc carbonate stops disappearing/dissolving <b>OR</b> a suspension (of zinc carbonate) forms <b>OR</b> the liquid turns cloudy	allow solid remains in the solution  ignore the reaction stops	1
(c)	filtration	accept filtering accept centrifuging	1
(d)	<b>M1</b> – <u>heat/boil</u> to <u>partially</u> evaporate (the water)  <b>M2</b> – leave to crystallise / leave until crystals form  <b>M3</b> – filter (to remove excess liquid)  <b>M4</b> – appropriate method of drying crystals	accept to remove <u>some</u> of the water accept heat to form a saturated/concentrated solution / heat until crystals form on (cold) glass rod / heat until crystals (just start to) form If evaporated to dryness then award no marks for whole question  accept leave to cool  accept pour off/decant (excess) liquid  e.g. use filter paper/blotting paper/kitchen towel / leave in (warm) oven/drying oven Accept leave to dry Do not accept hot oven/heat with a Bunsen flame	1  1  1

Question number	Answer	Notes	Marks
14 (a) (i)	<p><b>M1</b> – (covalent) bonds have to be broken</p> <p><b>M2</b> – large amount of energy required / bonds are strong</p> <p>(ii) the (covalent) bonding in silicon dioxide is stronger (than the (ionic) bonding in sodium chloride)</p>	<p>any mention of ions / metallic bonding / molecules / intermolecular forces scores 0/2</p> <p>Accept large number of bonds to be broken Accept forces (of attraction) between <u>atoms</u> in place of bonds</p> <p>Accept the covalent bonds (in silicon dioxide) are stronger than the ionic bonds (in sodium chloride) Accept more energy is required to break the (covalent) bonds in silicon dioxide (than is required to break the (ionic) bonds in sodium chloride) Accept forces (of attraction) between <u>atoms</u> in place of bonds</p>	1  1  1
(b)	<u>ions</u> flow/move (to the electrodes)	Accept ions are mobile/can move reject electrons	1
(c)	weak forces (of attraction) between <u>molecules</u> / weak <u>intermolecular</u> forces (of attraction) / little energy is required to separate <u>molecules</u>	Accept boiling point is below room temperature reject any mention of covalent bonds broken	1



Question number	Answer	Notes	Marks															
15 (a)	<table border="1" data-bbox="459 295 1485 443"> <tr> <td data-bbox="459 295 954 347">Titration number</td> <td data-bbox="954 295 1088 347">1</td> <td data-bbox="1088 295 1238 347">2</td> <td data-bbox="1238 295 1361 347">3</td> <td data-bbox="1361 295 1485 347">4</td> </tr> <tr> <td data-bbox="459 347 954 395">Volume of KMnO<sub>4</sub> solution added / cm<sup>3</sup></td> <td data-bbox="954 347 1088 395">22.80</td> <td data-bbox="1088 347 1238 395">22.10</td> <td data-bbox="1238 347 1361 395">22.50</td> <td data-bbox="1361 347 1485 395">22.20</td> </tr> <tr> <td data-bbox="459 395 954 443">Concordant titration results (✓)</td> <td data-bbox="954 395 1088 443"></td> <td data-bbox="1088 395 1238 443">✓</td> <td data-bbox="1238 395 1361 443"></td> <td data-bbox="1361 395 1485 443">✓</td> </tr> </table>	Titration number	1	2	3	4	Volume of KMnO <sub>4</sub> solution added / cm <sup>3</sup>	22.80	22.10	22.50	22.20	Concordant titration results (✓)		✓		✓		1
Titration number	1	2	3	4														
Volume of KMnO <sub>4</sub> solution added / cm <sup>3</sup>	22.80	22.10	22.50	22.20														
Concordant titration results (✓)		✓		✓														
(b)	<p data-bbox="459 539 795 614"><b>M1</b> <math>\frac{22.1(0) + 22.2(0)}{2}</math></p> <p data-bbox="459 778 712 817"><b>M2</b> – 22.15 (cm<sup>3</sup>)</p>	<p data-bbox="1518 539 1910 826">CSQ on boxes ticked in (a) If no results ticked, award <b>M1</b> only if columns 2 and 4 averaged If only one result ticked, no marks can be awarded in (b)</p> <p data-bbox="1518 865 1910 970">CSQ on results averaged, but the results must be taken from the table</p> <p data-bbox="1518 1008 1870 1152">Answer must be to 2dp correct answer with no working scores 2</p>	1  1															
(c)	D (pipette)		1															

Question number	Answer	Notes	Marks
15 (d) (i)	<p><b>M1</b> <math>\frac{20(.00) \times 0.02(00)}{1000}</math></p> <p>-</p> <p><b>M2</b> - <math>4(.00) \times 10^{-4}</math> (mol)</p>	0.4(00) scores 1	1
(ii)	$5 \times \mathbf{M2}$ from (i) / $4(.00) \times 10^{-4} \times 5 / 2(.00) \times 10^{-3}$		1
(iii)	$10 \times$ answer to (ii) / $2(.00) \times 10^{-2}$		1
(iv)	answer to (iii) $\times 152 / (2(.00) \times 10^{-2} \times 152) = 3.04$ (g)		1
(e) (i)	$m(\text{H}_2\text{O}) = (24.2 - 15.2) = 9(.0)$ (g)	must be given as a whole number	1
(ii)	answer to (i) $\div 18 / n(\text{H}_2\text{O}) = (9.00 \div 18) = 0.5(0)$ (mol)		1
(iii)	$n(\text{FeSO}_4) = (15.2 \div 152) = 0.1(00)$ (mol)		1
(iv)	$x =$ answer to (ii) $\div$ answer to (iii) / 5		1



Pearson Education Limited. Registered company number 872828  
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom